

Thermodynamics of Fateev's models in the Presence of External Fields

Davide Controzzi and Alexei M. Tsvelik

Department of Physics, University of Oxford, 1 Keble Road, Oxford, OX1 3NP, UK

(February 1, 2008)

We study the Thermodynamic Bethe Ansatz equations for a one-parameter quantum field theory recently introduced by V.A.Fateev. The presence of chemical potentials produces a kink condensate that modifies the excitation spectrum. For some combinations of the chemical potentials an additional gapless mode appears. Various energy scales emerge in the problem. An effective field theory, describing the low energy excitations, is also introduced.

PACS numbers: 11.10-z;71.10-w; 74.20De.

I. INTRODUCTION

Recently V.A.Fateev introduced a series of two dimensional integrable deformations of affine Toda theories possessing the property of duality^{1,2}. These models exist in two incarnations - fermionic and bosonic - and the strong coupling limit of one incarnation corresponds to the weak coupling limit of the other one. Apart from being amusing examples of integrable field theories, these models may have some interesting physical applications³⁻⁵.

In this paper we consider in some details one of the simplest models of that family. The Lagrangian density of the fermionic version of the theory is given by:

$$\mathcal{L}^{(f)} = \sum_{s=1,2} \left[i\bar{\psi}_s \gamma_\mu \partial_\mu \psi_s - \frac{\pi g}{2} (\bar{\psi}_s \gamma_\mu \psi_s)^2 \right] + \frac{1}{2} [(\partial_\mu \phi_1)^2 + (\partial_\mu \phi_2)^2] - M_0 \bar{\psi}_1 \psi_1 e^{-\beta \phi_1} - M_0 \bar{\psi}_2 \psi_2 e^{\beta \phi_2} - \frac{M_0^2}{2\beta^2} \left[2e^{\beta(\phi_2 - \phi_1)} + e^{-2\beta \phi_1} + e^{2\beta \phi_2} \right] \quad (1)$$

where $g = \beta^2/(4\pi + \beta^2)$ and we can identify two types of fermions $\psi_{1,2}$ and two coupled Toda chains described by the bosonic fields $\phi_{1,2}$.

In the bosonic representation the phonon modes, $\phi_{1,2}$, disappear and the Lagrangian density is :

$$\mathcal{L}^{(b)} = \frac{1}{2} \sum_{s=1}^2 \frac{\partial_\mu \chi_s \partial_\mu \bar{\chi}_s}{1 + (\gamma/2)^2 \chi_s \bar{\chi}_s} - \frac{M_0^2}{2} [\chi_1 \bar{\chi}_1 + \chi_2 \bar{\chi}_2 + \gamma^2/2 (\chi_1 \bar{\chi}_1)(\chi_2 \bar{\chi}_2)] \quad (2)$$

where the coupling constants of the two models are connected by the duality relation:

$$\gamma = 4\pi/\beta \quad (3)$$

As we have mentioned above, the weak coupling limit of the bosonic theory, $\gamma \ll 1$ ($1 - g \ll 1$), correspond to the strong coupling limit of the fermionic one, $\beta \gg 1$.

The bosonic form of the model is particularly interesting because the Euclidean version of the theory can be interpreted as Ginzburg-Landau free energy of two coupled superconductors. We shall discuss this application in greater details later in the text as well as in a separate publication.

Models (1),(2) have $U(1) \times U(1)$ symmetry and the corresponding conserved charges are given by

$$Q_s = \int dx \bar{\psi}_s \gamma_0 \psi_s = -\frac{i}{2} \int dx \frac{(\bar{\chi}_s \partial_0 \chi_s - \chi_s \partial_0 \bar{\chi}_s)}{[1 + (\gamma/2)^2 \chi_s \bar{\chi}_s]} \quad (4)$$

Thus one can introduce two chemical potentials and modify the Hamiltonian:

$$H \rightarrow H - h_1 Q_1 - h_2 Q_2 \quad (5)$$

The $U(1) \times U(1)$ symmetry of the Hamiltonian reflects in the symmetry of the particle multiplets. In absence of chemical potentials the spectrum of the model consists of fundamental particles carrying quantum numbers of $U(1)$ groups and their neutral bound states. The two-body scattering matrix of fundamental particles is given by a tensor product of two sine-Gordon S -matrices multiplied by a CDD-factor responsible for cancellation of double poles:

$$\mathcal{S}_{12}(\theta) = -\frac{\sinh \theta - i \sin(\pi/\lambda)}{\sinh \theta + i \sin(\pi/\lambda)} \hat{S}(\lambda; \theta) \times \hat{S}(\lambda; \theta) \quad (6)$$

where the $\hat{S}(\lambda, \theta)$ is the soliton scattering matrix of the sine-Gordon model with

$$\lambda = 2 - g = 1 + \frac{\gamma^2}{4\pi + \gamma^2} \quad (7)$$

and is given by

$$\hat{S}_{ab}^{\bar{a}, \bar{b}}(\lambda; \theta) = e^{i\delta_\lambda(\theta)} \left\{ \delta_{a\bar{a}} \delta_{b\bar{b}} \left[-\delta_{ab} + (1 - \delta_{ab}) \frac{\sinh(\lambda\theta)}{\sinh[\lambda(\theta - i\pi)]} \right] + \delta_{a\bar{b}} \delta_{b\bar{a}} (1 - \delta_{ab}) \frac{i \sin(\pi\lambda)}{\sinh[\lambda(\theta - i\pi)]} \right\} \quad (8)$$

with

$$\delta_\lambda(\theta) = \int_0^\infty d\omega \frac{\sin(\omega\theta)}{\omega} \frac{\sinh[\pi\omega(\lambda^{-1} - 1)/2]}{\cosh(\pi\omega/2) \sinh(\pi\omega/2\lambda)}$$

To derive thermodynamic equations we start from the discrete Bethe Ansatz equations which emerge when the system is put in a box of size L with periodic boundary conditions:

$$e^{-iM \sinh \theta_j L} \xi(a_1, \dots, a_n) = \prod_{k \neq j} \mathcal{S}(\theta_j - \theta_k) \xi(\bar{a}_1, \dots, \bar{a}_n) \quad (9)$$

$$E = M \sum_{j=1}^n \cosh \theta_j \quad (10)$$

where $\xi(a_1, \dots, a_n)$ is the wave function of n fundamental particles in the isotopic space. The matrix on the right-hand side of Eq.(9) is related to the trace of the monodromy matrix τ :

$$\prod_{k \neq j} \mathcal{S}(\theta_j - \theta_k) = \text{Tr} \tau(\lambda = \theta_j; \theta_1, \dots, \theta_n) \quad (11)$$

Up to the scalar factor given by the product of CDD-factors the latter matrix is equal to a tensor product of monodromy matrices of the sine-Gordon models:

$$\hat{\tau}(\lambda, \theta_1, \dots, \theta_n) = \hat{\tau}^{(1)}(\lambda, \theta_1, \dots, \theta_n) \times \hat{\tau}^{(2)}(\lambda, \theta_1, \dots, \theta_n) \quad (12)$$

The eigenvalues of the sine-Gordon monodromy matrix are known, hence it is straightforward to write down (9) in the diagonal form:

$$e^{-iM \sinh \theta_j L} = \prod_{k \neq j} S_0(\theta_j - \theta_k) \times \prod_{a=1}^{m_+} \frac{\sinh \lambda[(\theta_j - u_a) + i\pi/2]}{\sinh \lambda[(\theta_j - u_a) - i\pi/2]} \prod_{b=1}^{m_-} \frac{\sinh \lambda[(\theta_j - v_b) + i\pi/2]}{\sinh \lambda[(\theta_j - v_b) - i\pi/2]} \quad (13)$$

$$\prod_{j=1}^n \frac{\sinh \lambda[(\theta_j - u_a) + i\pi/2]}{\sinh \lambda[(\theta_j - u_a) - i\pi/2]} = \prod_{b=1}^{m_+} \frac{\sinh \lambda[(u_b - u_a) + i\pi]}{\sinh \lambda[(u_b - u_a) - i\pi]} \quad (14)$$

$$\prod_{j=1}^n \frac{\sinh \lambda[(\theta_j - v_a) + i\pi/2]}{\sinh \lambda[(\theta_j - v_a) - i\pi/2]} = \prod_{b=1}^{m_-} \frac{\sinh \lambda[(v_b - v_a) + i\pi]}{\sinh \lambda[(v_b - v_a) - i\pi]} \quad (15)$$

where:

$$S_0(\theta) = e^{2i\delta_\lambda(\theta)} \prod_{a=1}^{n-1} \frac{\sinh \theta - i \sin(\pi a/\lambda)}{\sinh \theta + i \sin(\pi a/\lambda)} \quad (16)$$

The numbers $m_\pm$ are related to the conserved charges $Q_{1,2}$. As we shall demonstrate later

$$m_\pm = \frac{1}{2} [n - (Q_1 \pm Q_2)] \quad (17)$$

The simplicity of this formula cannot obscure its remarkable meaning: propagating particles are superpositions of states localized on different edges of the stripe. This result is valid for other Fateev's models where the Toda array consists of more than two elastic chains.

II. THERMODYNAMIC BETHE ANSATZ (TBA) EQUATIONS

In order to construct the Thermodynamic Bethe Ansatz (TBA) equations from (13,14,15) we consider the case $1 - g = 1/\nu$, with integer $\nu \geq 2$, where the complex solutions of these equations (the strings) have the simplest classification. For most of the results the fact that ν is an integer is not important and one can generalize them for arbitrary $g > 1/2$ replacing ν by $(1 - g)^{-1}$.

To save space and make our notations more transparent we shall sometimes denote the kernels in the integral equations directly in terms of their Fourier transforms. For example, the convolution of two functions g and f with the function g having Fourier transform $g(\omega)$ will be written as:

$$\int_{-\infty}^{+\infty} d\theta' g(\theta - \theta') f(\theta') = g * f(\theta) = \{g(\omega)\} * f \quad (18)$$

Thermodynamic Bethe Ansatz (TBA) equations have been introduced in^{1,4}. The free energy of the system is expressed in terms of the excitation energies $E_n(\theta)$:

$$F/L = -T \sum_n M_n \int \frac{d\theta}{2\pi} \cosh \theta \ln[1 + e^{-E_n(\theta)/T}] \quad (19)$$

In the specific case (1),(2) we have only one bound state with energy E_1 , and the fundamental particle with energy E_0 . The function E_1 satisfies the following equation :

$$T \ln[1 + e^{E_1(\theta)/T}] - G_{11} * T \ln[1 + e^{-E_1(\theta)/T}] = M_1 \cosh \theta + G_{10} * T \ln[1 + e^{-E_0(\theta)/T}], \quad (20)$$

and is directly coupled only to E_0 , which is determined by the equation:

$$T \ln(1 + e^{E_0(\theta)/T}) - K_+ * T \ln[1 + e^{-E_0(\theta)/T}] = M \cosh \theta - \frac{h_+ + h_-}{2} + G_{10} * T \ln[1 + e^{-E_1(\theta)/T}] - Ts * \ln[(1 + e^{\epsilon_{-1}(\theta)/T})(1 + e^{\epsilon_1(\theta)/T})] \quad (21)$$

The two masses are related by:

$$M_1 = 2M \sin[\pi/2(2 - g)] \quad (22)$$

and the Fourier transform of the kernel is:

$$K_+ = K_{sym} + 2a_1 * s = 1 + \frac{\tanh[\pi\omega/2(\nu + 1)] \sinh[\pi(\nu - 1)\omega/2(\nu + 1)]}{\cosh(\pi\omega/2)} \quad (23)$$

where K_{sym} is related to the two-particle scattering phase :

$$K_{sym}(\theta) = \delta(\theta) - \frac{1}{2\pi i} \frac{d}{d\theta} \log(S_0(\theta)), \quad (24)$$

and its Fourier transform is:

$$K_{sym}(\omega) = \frac{\sinh \pi/2(1 - 1/\lambda)\omega \cosh(\pi\omega/\lambda)}{\cosh(\pi\omega/2) \sinh(\pi\omega/2\lambda)}. \quad (25)$$

The equations for ϵ_n and ϵ_{-n} are:

$$\epsilon_{-\nu} = h_{-}\nu/2 - s * T \ln[1 + e^{\epsilon_{-(\nu-2)/T}}] \quad (26)$$

$$\begin{aligned} \epsilon_{-n} = & \delta_{n,\nu-1} h_{-}\nu/2 + s * T \ln[1 + e^{-\epsilon_{n-1}/T}][1 + e^{\epsilon_{-n+1}/T}] \\ & + \delta_{\nu-2,n} s * T \ln[1 + e^{-\epsilon_{-\nu}/T}] + \delta_{n,1} s * T \ln[1 + e^{-E_0/T}] \end{aligned} \quad (27)$$

$$\epsilon_{\nu} = h_{+}\nu/2 - s * T \ln[1 + e^{\epsilon_{(\nu-2)/T}}] \quad (28)$$

$$\begin{aligned} \epsilon_n = & \delta_{n,\nu-1} h_{+}\nu/2 + s * T \ln[1 + e^{\epsilon_{n-1}/T}][1 + e^{\epsilon_{n+1}/T}] \\ & + \delta_{\nu-2,n} s * T \ln[1 + e^{-\epsilon_{\nu}/T}] + \delta_{n,1} s * T \ln[1 + e^{-E_0/T}] \end{aligned} \quad (29)$$

The Fourier transforms of the remaining kernels have the form:

$$s(\omega) = [2 \cosh[\pi/2\lambda(1 - g)\omega]]^{-1}$$

$$G_{11}(\omega) = \frac{4 \sinh(g\pi\omega/2\lambda) \sinh[\pi(1 - g)\omega/\lambda]}{\sinh(\pi\omega/2\lambda)}$$

$$\times \frac{\cosh[(\lambda - 1)\pi\omega/2\lambda] \sinh[\pi\omega/2\lambda]}{\cosh(\pi\omega/2)}$$

$$G_{10}(\omega) = \frac{2 \sinh(g\pi\omega/2\lambda) \sinh[\pi(1 - g)\omega/2\lambda]}{\cosh(\pi\omega/2)}$$

where λ is defined in (7) and can be also expressed as $\lambda = 1 + \nu^{-1}$.

The subsequent analysis will show that the fields $h_{\pm}$ are linear combinations of the chemical potentials $h_{1,2}$

$$h_{\pm} = h_1 \pm h_2 \quad (30)$$

(recall Eq.(17)).

III. PROPERTIES OF THE GROUND STATE

In this section we consider the zero temperature limit of the TBA equations. We assume that the fields $h_{\pm}$ are strong enough to make E_0 negative in some interval $[-B, B]$. Then it is obvious from Eq.(20) that $E_1 > 0$:

$$E_1 = M_1 \cosh \theta - G_{01} * E_0^{(-)} \quad (31)$$

From Eqs.(27,29) we deduce that all $\epsilon_{\pm n}$ with $n \neq \nu$ are also positive:

$$\epsilon_{\pm n} = -a_1 * E_0^{(-)} - A_{n,\nu-2} s * \epsilon_{\pm\nu}^{(-)} + (n - 1)h_{\pm} \quad (32)$$

where

$$\begin{aligned} A_{nm}(\omega) = & 2 \coth \Omega \frac{\sinh[(\nu - \max(n, m))\Omega] \sinh[\min(n, m)\Omega]}{\sinh \nu \Omega} \\ a_n(\Omega) = & \frac{\sinh[\pi(\nu - n)\Omega]}{\sinh \nu \Omega}, \quad \Omega = \frac{\pi\omega(1 - g)}{2(2 - g)} \end{aligned} \quad (33)$$

Substituting Eq.(32) into Eq.(26,28) we get the equations for $\epsilon_{\pm\nu}$:

$$\epsilon_{\pm\nu}^{(+)} + \left\{ 1 - \frac{\sinh[(\nu-2)\Omega]}{\sinh \nu\Omega} \right\} * \epsilon_{\pm\nu}^{(-)} = h_{\pm} + \left\{ \frac{\sinh \Omega}{\sinh \nu\Omega} \right\} * E_0^{(-)} \quad (34)$$

At last, substituting Eq.(32) into Eq.(21) we get:

$$E_0^{(+)} + K_{sym} * E_0^{(-)} = M \cosh \theta - \frac{1}{2}(h_+ + h_-) + \left\{ \frac{\sinh \Omega}{\sinh \nu\Omega} \right\} * [\epsilon_{\nu}^{(-)} + \epsilon_{-\nu}^{(-)}]. \quad (35)$$

Eqs.(34,35) determine the structure of the ground state. We cannot solve them in the general case and therefore we shall consider only two special cases: a) $h_+ \ll h_-$; b) $h_+ \approx h_-$.

In the first case $\epsilon_{-\nu} > 0$ and ϵ_{ν} is negative inside an interval $[-Q, Q]$, with $Q \gg B$ and $Q \rightarrow \infty$ as $h_+ \rightarrow 0$. Since $\epsilon_{\nu}^{(+)}$ is small it is convenient to invert the kernel in Eq.(34) to get:

$$\left\{ \frac{\sinh \nu\Omega}{2 \sinh \Omega \cosh[(\nu-1)\Omega]} \right\} * \epsilon_{\nu}^{(+)} + \epsilon_{\nu}^{(-)} = \nu h_+/2 + \left\{ \frac{1}{2 \cosh[(\nu-1)\Omega]} \right\} * E_0^{(-)} \quad (36)$$

For $|\theta| \gg B$ it is useful to approximate the right hand side of this equation by its asymptotics:

$$\left\{ \frac{1}{2 \cosh[(\nu-1)\Omega]} \right\} * E_0^{(-)} \simeq \frac{2\zeta}{\pi} a_{\epsilon} \exp(-\zeta|\theta|) \quad (37)$$

where

$$a_{\epsilon} = \int_{-B}^B d\theta' E_0(\theta') e^{-\zeta\theta'} \quad (38)$$

and

$$\zeta = \frac{\nu+1}{\nu-1} = \frac{2-g}{g} \quad (39)$$




FIG. 1. Schematic diagram of the ground state energies in presence of chemical potential for $h_+ \ll h_-$ and large B. The same situation for small B will be presented in section VII

Analogously one can rewrite Eq.(35) as:

$$E_0^{(+)} + K * E_0^{(-)} = M \cosh \theta - h_-/2 - \left\{ \frac{1}{2 \cosh[(\nu-1)\Omega]} \right\} * \epsilon_{\nu}^{(+)} \quad (40)$$

where

$$K(\Omega) = \frac{\sinh \Omega \cosh[(3\nu - 1)\Omega]}{2 \sinh \nu \Omega \cosh[(\nu - 1)\Omega] \cosh[(\nu + 1)\Omega]} \quad (41)$$

Since $\epsilon_\nu^{(+)}$ is very small, to first approximation this reduces to:

$$E_0^{(+)} + K * E_0^{(-)} = M \cosh \theta - h_-/2 \quad (42)$$

For large B we can approximate Eqs.(36) and (42) by the Wiener-Hopf (WH) ones⁷. The solutions for $\epsilon_0^{(-)}(\theta) = E_0^{(-)}(\theta + B)$ and $\epsilon_\nu^{(+)}(\theta) = \epsilon_\nu^{(+)}(\theta + B)$ are:

$$\epsilon_0^{(-)}(\omega) = \frac{h_-}{G_a^{(-)}(\omega)G_a^{(+)}(0)} \left[\frac{1}{i\omega + 1} - \frac{1}{i\omega + 0} \right] \quad (43)$$

$$\epsilon_\nu^{(+)}(\omega) = \frac{1}{G^{(+)}(\omega)} \left[\frac{h_+}{G^{(-)}(0) - i\omega + 0} - a_\epsilon \frac{e^{-\zeta B}}{G^{(-)}(i\zeta)} \frac{1}{\zeta - i\omega} \right] \quad (44)$$

where a_ϵ is defined in Eq.(38), and $G_a^{(+)}(G_a^{(-)})$ and $G^{(+)}(G^{(-)})$ are analytic in the upper (lower) half plane and satisfy the conditions:

$$\begin{aligned} G_a^{(+)}(\omega)G_a^{(-)}(\omega) &= K(\omega) \\ G^{(+)}(\omega)G^{(-)}(\omega) &= \frac{\sinh \nu \theta}{2 \sinh \Omega \cosh[(\nu - 1)\Omega]}; \end{aligned} \quad (45)$$

ζ is defined by Eq.(39) and B is determined by the following relation:

$$\frac{Me^B}{G^{(+)}(i)} = \frac{h_-}{G^{(-)}(0)} \quad (46)$$

To study the low temperature thermodynamic in Section VI it is useful to introduce also the ground state densities for E_0 and ϵ_ν , σ and Σ respectively, defined by the following equations:

$$\sigma^{(+)} + K * \sigma^{(-)} = \frac{M}{2\pi} \cosh \theta \quad (47)$$

$$\Sigma^{(-)} + \left\{ \frac{\sinh \nu \Omega}{2 \sinh \Omega \cosh[(\nu - 1)\Omega]} \right\} * \Sigma^{(+)} = \left\{ \frac{1}{2 \cosh[(\nu - 1)\Omega]} \right\} * \sigma^{(-)} \equiv \Sigma^0 \quad (48)$$

Solutions of the WH equations for $\tilde{\sigma}^{(-)}(\theta) = \sigma^{(-)}(\theta + B)$ and $\tilde{\Sigma}(\theta) = \Sigma(\theta + B)$ are:

$$\tilde{\sigma}^{(-)}(\omega) = \frac{h_-}{4\pi G^{(-)}(\omega)G^{(+)}(i)(i\omega + 1)} \quad (49)$$

$$\tilde{\Sigma}(\omega) = \frac{1}{G^{(+)}(\omega)} \left[\frac{h_+}{G^{(-)}(0)(-i\omega + 0)} - a_\epsilon \frac{e^{-\zeta B}}{G^{(-)}(i\zeta)(\zeta - i\omega)} \right] \quad (50)$$

In the limit b) both $\epsilon_{\pm\nu}$ are positive and we have for E_0 :

$$E_0^{(+)} + K_{sym} * E_0^{(-)} = M \cosh \theta - h \quad (51)$$

Again using the WH method we find the following solution:

$$\epsilon_0^{(-)} = \frac{h}{G_b^{(-)}(\omega)G_b^{(+)}(0)(i\omega + 1)(i\omega + \delta)} \quad (52)$$

with $G_b^{(+)}(\omega)G_b^{(-)}(\omega) = K_{sym}(\omega)$. Substituting this solution in Eq.(34) we obtain the following estimate for the gap in $\epsilon_{\pm\nu}$:

$$M_\nu \sim h(M/h)^{(2-g)/2} \quad (53)$$

In the limit $\nu^{-1} \rightarrow 0 (g \rightarrow 1) M_\nu \sim \sqrt{Mh}$.

A schematic diagram of these results is presented in Fig.1 and Fig.2 for the case a) and b) respectively.




FIG. 2. Schematic diagram of the ground state energies in presence of chemical potential for $h_+ \approx h_-$ and large B.

IV. THE LIMIT OF FREE BOSONS

To check the validity of TBA equations (20-29) and establish the connection between the fields $h_\pm$ and the chemical potentials $h_{1,2}$ we consider the free boson limit of the TBAs. It is realized in the case $g \rightarrow 1$ or $\nu^{-1} \rightarrow 0$. In this case the kernels in Eqs.(26-29) become proportional to the delta functions

$$s(\theta) \rightarrow \frac{1}{2}\delta(\theta)$$

and $K_+ \rightarrow I, G_{11} = G_{01} \rightarrow 0$, such that all TBA equations become algebraic. The mass of the bound state in this limit is $M_1 = 2M$ and the Free Energy is equal to:

$$F/L = -\frac{TM}{2\pi} \int d\theta \cosh \theta \ln \left\{ \left[1 + e^{-E_1/T} \right]^2 \left[1 + e^{-E_0/T} \right] \right\} \quad (54)$$

The solution of TBA equations is expressed in terms of $\xi = M \cosh \theta/T$:

$$1 + e^{-E_1/T} = (1 - e^{-2\xi})^{-1} \quad (55)$$

$$1 + e^{\epsilon_{\pm n}/T} = \left[\frac{\sinh(h_\pm n + a_\pm)/2T}{\sinh h_\pm/2T} \right]^2 \quad (56)$$

where

$$\frac{\sinh a_\pm/2T}{\sinh h_\pm/2T} = \sqrt{1 + e^{-E_0/T}} \quad (57)$$

Substituting this into Eq.(21) we get

$$\left\{ \cosh(h_+/2T) \sqrt{1 + e^{-E_0/T}} + [\cosh^2(h_+/2T) + \sinh^2(h_+/2T) e^{-E_0/T}]^{1/2} \right\} \times$$

$$\left\{ \cosh(h_-/2T) \sqrt{1 + e^{-E_0/T}} + [\cosh^2(h_-/2T) + \sinh^2(h_-/2T) e^{-E_0/T}]^{1/2} \right\} = e^{\xi - E_0/T} \quad (58)$$

which has the following solution:

$$1 + e^{-E_0/T} = \frac{\sinh^2 \xi}{(\cosh \xi - \cosh(h_+/2T) \cosh(h_-/2T))^2 - \sinh(h_+/2T) \sinh(h_-/2T)} \quad (59)$$

Combining it with (55) in the Free Energy, we get

$$F/L = -\frac{T}{2\pi} \int d\theta M \cosh \theta \ln(\mathcal{F}) \quad (60)$$

$$\begin{aligned} \mathcal{F} &= (1 - e^{-2\xi})^{-2} \frac{\sinh^2 \xi}{(\cosh \xi - \cosh(h_+/2T) \cosh(h_-/2T))^2 - \sinh^2(h_+/2T) \sinh^2(h_-/2T)} \\ &= \left\{ [1 - e^{-\xi - (h_+ + h_-)/2T}] [1 - e^{-\xi + (h_+ + h_-)/2T}] [1 - e^{-\xi - (h_+ - h_-)/2T}] [1 - e^{-\xi + (h_+ - h_-)/2T}] \right\}^{-1} \end{aligned} \quad (61)$$

The free boson limit of TBAs (20-29) does reproduce the free energy of two non interacting complex bosonic fields. This indicates that the TBA equations are correct. Another important result is the confirmation of Eqs.(17,30).

V. SEMICLASSICAL ANALYSIS OF THE LIMIT OF SMALL γ

Let us consider the Euclidean form of the Lagrangian (2). Rescaling the fields, $\chi_s \rightarrow (\gamma/2)\Delta_s$, it becomes:

$$\mathcal{L}^{(b)} = \frac{2}{\gamma^2} \left[\sum_{s=1}^2 \frac{\partial_\mu \Delta_s \partial_\mu \bar{\Delta}_s}{1 + \Delta_s \bar{\Delta}_s} + M^2 \{ \Delta_1 \bar{\Delta}_1 + \Delta_2 \bar{\Delta}_2 + 2(\Delta_1 \bar{\Delta}_1)(\Delta_2 \bar{\Delta}_2) \} \right] \quad (62)$$

Then in the limit $\gamma \rightarrow 0$ the quantum fluctuations are suppressed and one can approximate the ground state energy as minimum of the functional

$$E(\Delta_s, h_s) = \frac{2}{\gamma^2} \left[M^2 \{ \Delta_1 \bar{\Delta}_1 + \Delta_2 \bar{\Delta}_2 + 2(\Delta_1 \bar{\Delta}_1)(\Delta_2 \bar{\Delta}_2) \} - \sum_{s=1,2} \frac{h_s \Delta_s \bar{\Delta}_s}{1 + \Delta_s \bar{\Delta}_s} \right]. \quad (63)$$

We will not address this problem, already studied by Fateev¹, but we will use this approximation to get more insight to the problem and give an intuitive picture of the low energy physics. Let us perform the following transformation in the form (62):

$$\Delta_s = \sinh \rho_s e^{i\varphi_s} \quad (64)$$

Under this transformation the measure transform in $D\rho D\varphi \sinh(2\varphi)$ and the Lagrangian density in the new fields in presence of chemical potentials assume the form:

$$\begin{aligned} \mathcal{L}^{(b)} &= \frac{2}{\gamma^2} \left\{ \sum_{s=1,2} [(\partial_\mu \rho_s)^2 + \tanh^2 \rho_s (\partial_\mu \varphi_s)^2 - h_s \tanh^2 \rho_s \partial_0 \varphi_s] \right. \\ &\quad \left. + M^2 [\sinh^2 \rho_1 + \sinh^2 \rho_2 + 2 \sinh^2 \rho_1 \sinh^2 \rho_2] \right\} \end{aligned} \quad (65)$$

For $h_1 = h$ and $h_2 = 0$, which corresponds to the situation b in section III, we get:

$$\begin{aligned} \mathcal{L}^{(b)} &= \frac{2}{\gamma^2} \{ (\partial_\mu \rho_1)^2 + \tanh^2 \rho_1 [(\partial_\mu \varphi_1)^2 - h \partial_0 \varphi_1] + (\partial_\mu \rho_2)^2 + \\ &\quad \tanh^2 \rho_2 (\partial_\mu \varphi_2)^2 + M^2 (\sinh^2 \rho_1 + \sinh^2 \rho_2 + 2 \sinh^2 \rho_1 \sinh^2 \rho_2) \} \end{aligned} \quad (66)$$

The term $h(\partial_0 \varphi_1)$ can be absorbed by the shift:

$$\varphi_1 \rightarrow \varphi_1 - (h/2)t \quad (67)$$

which generate also an additional term $-h^2/4$. Then the Lagrangian density becomes:

$$\mathcal{L}^{(b)} = \frac{2}{\gamma^2} \{ (\partial_\mu \varphi_1)^2 + (\partial_\mu \varphi_2)^2 + \tanh^2 \rho_1 (\partial_\mu \varphi_1)^2 + \tanh^2 \rho_2 (\partial_\mu \varphi_2)^2 + V_{eff}^b(\rho_1, \rho_2) \} \quad (68)$$

where:

$$V_{eff}^b(\rho_1, \rho_2) = V_1(\rho_1) + M^2 \sinh^2 \rho_2 + 2M^2 \sinh^2 \rho_1 \sinh^2 \rho_2 \quad (69)$$

and

$$V_1(\rho_1) = M^2 \sinh^2 \rho_1 - h^2/4 \tanh^2 \rho_1 \quad (70)$$

For $h/2 > M$, $V_{eff}^b(\rho_1, \rho_2)$ has a minimum at $\rho_1 = \rho_0$ and $\rho_2 = 0$ where $\exp 2\rho_0 = \sqrt{8}(h/M)$. In the vicinity of the minimum we have:

$$V_{eff}(\rho_1 = \rho_0 + x, \rho_2 = y) \approx hM(\sqrt{2}y^2 + \sqrt{8}x^2) \quad (71)$$

which gives the masses going like $\sqrt{hM}$, in accordance with (53).

Thus when the chemical potential exceeds the threshold the kinks condense and gapless Goldstone mode appear. To see this explicitly we write $\rho_1 = \rho_0 + x$ and keeping only quadratic terms in x obtain the following expression for the Lagrangian:

$$\begin{aligned} \mathcal{L}^{(b)} \sim \frac{2}{\gamma^2} \{ (\partial_\mu x)^2 + (Mh/2)x^2 + \tanh^2 \rho_0 (\partial_\mu \varphi_1)^2 + \\ (\partial_\mu \rho_2)^2 + \tanh^2 \rho_2 (\partial_\mu \varphi_2)^2 + \sinh^2 \rho_2 h/M \} \end{aligned} \quad (72)$$

Here we can identify a gapless mode φ_1 with velocity equal to the bare one, a gapful field x with the mass $m_x^2 = hM/2$ and an effective integrable field theory described by the Lagrangian density:

$$\mathcal{L}_2 = \frac{1}{2} \left(\frac{2}{\gamma} \right)^2 \{ M \sinh^2 \rho_2 + (\partial_\mu \rho_2)^2 + \tanh^2 \rho_2 (\partial_\mu \varphi_2)^2 \} \quad (73)$$

In the case $h_1 \sim h_2$ corresponding to the situation a) of Section III, the effective potential acquires the form:

$$V_{eff}^a(\rho_1, \rho_2) = V_1(\rho_1) + V_1(\rho_2) + 2M^2 \sinh^2 \rho_1 \sinh^2 \rho_2 \quad (74)$$

Repeating the same analysis one finds two independent gapless modes with the same velocity. Then the semiclassical approximation gives a correct qualitative description of the low energy behavior of the system. As a byproduct we obtain another confirmation of (30).

VI. LOW ENERGY PHYSICS: MASSLESS MODES AND ENERGY SCALES

Let us go back to the zero temperature limit described in Section III. As we have shown, in the case $h_1 \approx h_2$ there are two soft modes: E_0 and ϵ_ν . At $|\theta| < B$ the mode ϵ_ν closely follows E_0 ; at $|\theta| \gg B$ its behavior is determined by the asymptotics (37) and this mode becomes independent. The temperature scale below which the two modes decouple can be determined by the value of ϵ_ν at $\theta \sim B$:

$$T_{sep} \sim \epsilon_\nu^{(-)}(B) \quad (75)$$

For low temperatures, $T < T_{sep}$, the two modes give independent contribution to the thermodynamics. To calculate the low temperature free energy:

$$F/L = -\frac{T}{2\pi} M \int d\theta \cosh \theta \ln[1 + e^{-E_0(\theta)/T}] \quad (76)$$

we use the Eqs. (36,42) which at finite temperature have the form:

$$T \ln(1 + e^{E_0/T}) - K * T \ln(1 + e^{-E_0/T}) = M \cosh \theta - \frac{h_-}{2} - \left\{ \frac{1}{2 \cosh(\nu - 1)\Omega} \right\} * T \ln(1 + e^{\epsilon_\nu/T}) \quad (77)$$

$$\left\{ \frac{\sinh \nu \Omega}{2 \sinh \Omega \cosh(1 - \nu)\Omega} \right\} * T \ln(1 + e^{\epsilon_\nu/T}) - T \ln(1 + e^{-\epsilon_\nu/T}) = \frac{\nu h_+}{2} - a_\epsilon \quad (78)$$

where a_ϵ is defined in Eq.(37). To isolate the contributions that vanish at $T=0$ we rewrite (77) like:

$$E_0^{(+)} + K * E_0^{(-)} + (I - K) * T \ln(1 + e^{-|E_0|/T}) = \text{r.h.s. of Eq.(77)} \quad (79)$$

where I is the identity operator. Using Eq.(47) we can rewrite the free energy as:

$$F/L = -\frac{T}{2\pi} M \int d\theta \cosh \theta (E_0^{(-)} + T \ln(1 + e^{-|E_0|/T})) \equiv f_0 + f_\nu \quad (80)$$

where

$$f_0 = -T \int d\theta \sigma(\theta) \ln(1 + e^{-|E_0|/T}) \approx -\pi^2 T^2 / 3V_c \quad (81)$$

$$f_\nu = -T \int d\theta \left\{ \frac{1}{2 \cosh(\nu - 1)\Omega} \right\} \sigma^{(-)}(\theta) T \ln(1 + e^{\epsilon_\nu/T}) \approx -\pi^2 T^2 / 3V_s \quad (82)$$

The velocities of the two modes are determined by :

$$V_c = \sum_{s=\pm} \left| \frac{\partial E_0(\theta)}{\partial \theta} \right| [2\pi \sigma^{(s)}(B)]_{\theta=sB}^{-1} \quad (83)$$

$$V_s = \sum_{s=\pm} \left| \frac{\partial \epsilon_\nu(\theta)}{\partial \theta} \right| [2\pi \Sigma^{(s)}(B)]_{\theta=sB}^{-1} \quad (84)$$

For large B all the quantities appearing in these equations can be calculated using the Wiener-Hopf solutions obtained in Section III via the following relations:

$$\begin{aligned} \left| dE_0^{(\pm)} / d\theta \right|_{\theta=B} &= \left| d\epsilon_0^{(\pm)} / d\theta \right|_{\theta=0} = \lim_{\omega \rightarrow \infty} \pm (i\omega)^2 \epsilon^{(\pm)}(\omega) \\ \sigma^{(\pm)}(B) &= \tilde{\sigma}^{(\pm)}(0) = \lim_{\omega \rightarrow \infty} \pm i\omega \tilde{\sigma}^{(+)}(\omega) \end{aligned}$$

and analogous relations for ϵ_ν and Σ . The results are:

$$V_c = V_s = 1 \quad (85)$$

The fact that the two velocities are the same seems to be peculiar to the model in consideration and not a general property of the family of Fateev's models. In general V_s depends on g and on the number of Toda chains.

Thus we have shown that the presence of chemical potentials introduces additional energy scales in the problem with non trivial crossovers. At temperatures $T \ll h_-$, ϵ_{-n} and ϵ_n ($n \neq \nu$) modes decouple from the other modes. The same happens for E_1 at energies smaller than their gap and in some physical regions there is also an intermediate energy scale, as we will see in the next Section. As one lowers the temperature below T_{sep} the gapless modes decouple from each other and the picture of two non-interacting Luttinger liquids emerges.

VII. WEAK COUPLING LIMIT: NON PERTURBATIVE RESULTS

Now we wish to focus our attention on the limit $g \rightarrow 1$ which corresponds to the weak-coupling limit of the bosonic theory. Naively one can imagine to obtain perturbative results. It turns out however that, in the presence of the chemical potential, this is not the case, the reason being that the kinks condensate survives and affects the excitation spectrum. A similar effect is present in the Klein-Gordon limit of the sine-Gordon model⁶.

To be specific let us consider the situation a) of Section III, all the energies ϵ_n ($n \neq \nu$) have gaps of the order of h_- and can be omitted at temperatures $T \ll h_-$. To obtain information on the excitation spectrum one has to study the fundamental equations (31) and (42). The standard relativistic dispersion law: $E_1(\theta) = M_1 \cosh \theta$, can be modified by the last term in (31) which emerge in presence of kinks.

In the limit $g \rightarrow 1$, the kernels simplify $K(\omega) = K_0(\omega)(1 + O((1 - g)^2))$, and $G_{10}(\omega) = G_0(\omega)(1 + O((1 - g)^2))$, where

$$\begin{aligned} K_0(\omega) &= \frac{4\pi^2}{\beta^2\lambda} \omega \frac{\cosh(\pi\omega(\lambda + 2)/2\lambda)}{\cosh(\pi\omega/2) \sinh(\pi\omega/\lambda)} \\ &= \frac{\pi}{\lambda} (1 - g) \omega \frac{\cosh(\pi\omega(\lambda + 2)/2\lambda)}{\cosh(\pi\omega/2) \sinh(\pi\omega/\lambda)} \end{aligned} \quad (86)$$

and

$$G_0 = (1 - g) \frac{\pi\omega}{\lambda} \frac{\sinh(\pi\omega/2\lambda)}{\cosh(\pi\omega/2)} \quad (87)$$

Again we can study TBA equations with the kernel $K_0(\omega)$ and $G_0(\omega)$ in the limits of large and small B . We focus the attention on this second case where the effects we want to consider can be seen more clearly. To study Eq.(31) for small B we need the kernel $K_0(\theta)$ only for small θ . This is equivalent to approximating Eq.(86) for large ω , as:

$$K_0(\omega) \simeq \frac{2\pi}{\lambda} (1 - g) |\omega| \quad (88)$$

Approximating also the rhs of Eq. (42) for small θ we can rewrite it as:

$$K_0 \tilde{*} E_0 = M\theta^2/2 - H \quad (89)$$

where $H = h_-/2 - M$ and $\tilde{*}$ denote the convolution of support $(-B, B)$ and K_0 is the Fourier transform of (88). One can absorb the constant part of the kernel in E_0 introducing the new function $\tilde{E}_0 = \frac{2\pi}{\lambda} (1 - g) E_0$. Equation (42) then becomes:

$$\tilde{K}_0 \tilde{*} \tilde{E}_0 = M\theta^2/2 - H, \quad (90)$$

$\tilde{K}_0$ is the singular integral operator corresponding to the kernel $\tilde{K}_0(\omega) = |\omega|$ and defined:

$$\tilde{K}_0 * \tilde{E}_0(\theta) = -\frac{1}{\pi} P \int_{-B}^B \frac{1}{(\theta - \theta')^2} \tilde{E}_0(\theta') d\theta' = \frac{1}{\pi} \frac{\partial}{\partial \theta} P \int_{-B}^B \frac{1}{(\theta - \theta')} \tilde{E}_0(\theta') d\theta' \quad (91)$$




FIG. 3. Dispersion relation E_1 (upper figure) in the presence of kinks for $B \ll 1$ and $g = 0.02$, from the numerical solution of Eq.(31). The dotted line represents the solution in the absence of kink condensate. In the lower figure we plot the gapless mode E_0 .

Using this form of the kernel, Eq. (90) can be reduced to the canonical form:

$$P \int_{-B}^B d\theta' \frac{1}{(\theta - \theta')} V(\theta') = g(\theta) \quad (92)$$

where we have introduced: $V(\theta) = \partial \tilde{E}_0(\theta) / \partial \theta$ and $g(\theta) = \pi(M\theta^2/2 - H)$. This equation has the following general solution⁸:

$$V(\theta) = \frac{1}{\pi^2} \sqrt{B^2 - \theta^2} P \int_{-B}^B d\theta' \frac{g(\theta')}{\sqrt{B^2 - \theta'^2}(\theta' - \theta)} \quad (93)$$

which gives:

$$E_0(\theta) = -\frac{M\lambda}{3\pi^2(1-g)}(B^2 - \theta^2)^{3/2} \quad (94)$$

where $B^2/4 = \frac{h_-/2 - M}{M} \equiv \Delta$. Analogously the ground state density (47) turns out to be:

$$\sigma(\theta) = \frac{M\lambda}{2\pi^3(1-g)} \sqrt{B^2 - \theta^2} \quad (95)$$

From here we find the relationship between h_- and the charge:

$$Q = \frac{h_-/2 - M}{2\pi(1-g)} \quad (96)$$

The Fermi energy, $E_F = E_0(\theta)$ as function of the charge if therefore:

$$E_F = \frac{8M}{3\pi^2}(1-g)^{1/2} \left[\frac{2\pi}{M} Q \right]^{3/2} \quad (97)$$

We can now use the above results to solve Eq(31). Rewriting the Fourier transform of the kernel G_0 as:

$$G_0 \sim \frac{2}{\pi} \frac{\partial}{\partial \theta} \frac{(\pi/2(1-g)) \sinh \theta}{\sinh^2 \theta + (\pi/2(1-g))^2 \cosh^2 \theta} \quad (98)$$

it is possible to extract the asymptotic behavior of E_1 :

$$E_1 = \begin{cases} M_1 \cosh \theta + M/3 \cos(2 \sin^{-1}(\theta/B)), \theta \ll B \\ M_1 \cosh \theta + A e^{-\theta}, \theta \gg B \end{cases} \quad (99)$$

The numerical solution is shown on Fig. 3. One can clearly see that even in the weak coupling limit the dispersion law is drastically modified at small θ .

This analysis is supposed to clarify the subject of different energy scales present in the model (cfr.Fig.4). At very small doping, $Q \ll Q_1 \equiv M(g-1)^{-1/3}$, E_F is smaller than M and there are two scales besides h : M and E_F itself. The mode E_1 is completely decoupled at low temperature. As you increase the doping you reach an intermediate region, characterized by $M(1-g)^{-1/3} \ll Q \ll Q_2 \equiv M(1-g)^{-1}$, where $E_F \simeq M$ but still much smaller than E_1 . At very large doping we find a very interesting region. As we have seen in Sec. III at large Q (or equivalently at large B) $E_F \simeq Q$, while $E_1(0) \simeq \sqrt{Q}$. Then there is a regime in which E_F is bigger than the gap of the E_1 -mode, and there will be temperatures, $T \ll E_F$ for which the E_1 -mode becomes soft.




FIG. 4. A Schematic diagram describing different energy scales as a function of the charge. We can recognize three distinct regions as described in Sec. VII.

VIII. DISCUSSION

We have considered in some detail the thermodynamics of one of the integrable field theories recently introduced by Fateev. Most of the results are quite general and remain valid also for other models of the family. These models are interesting for various reasons. First of all they have peculiar mathematical properties mostly related to a dual representation of the theory. In addition they probably can find application to various physical systems. Considering the model as a (1+1)-dimensional Quantum Field Theory one can interpret it as two one dimensional conductors coupled via phonon interaction. The problem of quasi one dimensional systems with electron-phonon interaction is a very important one and cannot be approached with perturbative methods, it is then very important to have exact results for some specific model. This interpretation of the model and the possible application to physical systems have been discussed in ref.⁴, for physical regimes different from the one considered here (i.e. for $g < 1/2$). On the other side, the Euclidean version of the model, in the bosonic representation, describes an effective Landau-Ginzburg theory of two coupled superconductors. In the specific model considered in this work, the coupling between the superconductors is quite simple. Nevertheless for other models of the Fateev's family, where coupling between the fermionic modes is achieved via a higher number of Toda chains, the resulting coupling between the two superconductors is via elastic modes and then can be interpreted as two layered superconductors separated by an insulating stratum. In relation to the high- T_c (cuprate) superconductors an interesting problem is to study the dependence of the superconducting properties on the number of insulating layers between the superconducting planes. We will address this problem in a separate publication.

Let us give a brief summary of our results. As shown in Sec.VII the results are non-perturbative also in the weak coupling regime and could only be obtained through exact or non-perturbative methods. This is related to the fact that the chemical potentials generate a kink condensate that survives in the weak coupling limit. As we have shown, for some combinations of the chemical potentials two gapless modes emerge and various energy scales appears, which characterizes crossovers between different regimes. Probably the most interesting region of the phase diagram is obtained at large doping where the E_1 -mode become soft at temperatures for which the E_0 -one is still frozen. It deserves further investigation. Another important result is the one contained in Eq.(17). This equation tells us that charged particles in the Fateev's model are linear combinations of particles located on the edges, which is important for qualitative understanding of the physics of the model.

- ¹ V. A. Fateev, Nucl. Phys. B**479**, 594 (1996).
- ² V. A. Fateev, Nucl. Phys. B**473**, 509 (1996).
- ³ C. Pepin and A. M. Tsvetik, Phys. Rev. Lett. , **82**, 3859 (1999).
- ⁴ C. Pepin and A. M. Tsvetik, cond-mat 9983180.
- ⁵ D. Controzzi and A. M. Tsvetik, unpublished.
- ⁶ J.-S. Caux and A. M. Tsvetik, Nucl.Phys.B **474**, 715 (1996).
- ⁷ V.A.Fateev, E.Onofri, Al.A.Zamolodchikov, Nucl.Phys. B**406**,521 (1993).
- ⁸ S.G.Mikhlin, Integral equations (Pergmon press, Oxford,1964).